

5  
Physicochemical study of oil fractions from crude oil. I.  
Method and experimental technique. V. Kelio, V. Palo,  
and A. Tkáč (Slovenská vysoká škola, Bratislava,  
Czech.). *Chem. zvesti* 13, 205-211 (1959) (German summary).  
—Lubricating oil distillates from Gbely (Slovakia) crude oil  
were evaluated as raw materials for cable oil by means of  
the *n*-d.-mol. wt. method, elementary analysis, viscosity and  
viscosity index, dielec. losses and dielec. const., insulating  
properties, and infrared absorption spectroscopy. All tests  
were applied to chromatographic fractions of the distillates.  
For kinetic studies by infrared spectroscopy, oxidation of a  
thin hydrocarbon layer (1 mm.) was used. Jan Miska

KELLO, V.

Distr: 4E2c(j)/4E3b

Mean life of free radicals in solid natural rubber. A. Tkáč and V. Kello (Slovakian Coll. Technol., Bratislava, Czech.). *Trans. Faraday Soc.* 55, 1211-20 (1959); cf. C.A. 49, 6639k; 50, 8317i, 16337d. Free radicals were generated in solid films of natural rubber 20-30  $\mu$  thick on NaCl plates when heated in an autoclave for  $\sim 1000$  min. at 140°. The mean life of the free radical was detd. by a kinetic analysis of the subsequent oxidn. reactions. Radicals of relatively long life formed only in the presence of natural antioxidants. In the absence of antioxidants, heat-treatment reduced the sensitivity towards oxidn., a result which was explained by cross-linking with rapid removal of the radicals.

Victor R. Deltz

5  
1-28 (NB)  
2

KRCMERY, Vladimír; FERENCIK, Miroslav; KELLO, Vojtech

Activity of the hydroperoxydases in brucellosis. Biologia 14 no.12:  
924-932 '59. (EEAI 9:7)

1. Statny vedecky veterinarni ustav, Bratislava a Katedra  
fyzikalnej chemie Slovenskej vysokej školy technickej, Bratislava  
(BRUCELOSIS)  
(HYDROPEROXIDASES)

TKAC, Alexander, doc., dr., inz.; KELLO, Vojtech, prof., dr.

Problem of polyisoprene stability from the viewpoint of radical processes. Chem zvesti 17 no.4:237-247 '63.

1. Vedecky ustav pri Katedre fyzikalnej chemie, Slovenska vysoka skola technicka, Bratislava, Kollarovo namesti 2.

CZECHOSLOVAKIA

TKAC, A.; KELLO, V.; HRIVIKOVA, J.

Dept. of Physical Chemistry, Slovak Technical Univ., Bratislava (for all)

Prague, Collection of Czechoslovak Chemical Communications, No 2, Feb  
1966, pp 551-565

"On the theory of macroradical termination. Part 4: Mechanism of  
termination of macroradicals."

KHASYSKI, M.; KEL'M, M. [Kelm, M.]; SHIDLOVSKAYA, S. [Szydłowska, S.];  
GORAK, B. [Horak, B.]; RIKHTER, V.

From public reports of the heads of the delegations of socialist  
countries. Tekh. est. 2 no.8:7-11 Ag '65. (MIRA 18:9)

1. Direktor TSentra promyshlennoy estetiki, khudozhestvennogo  
proyektirovaniya i konstruirovaniya Narodnoy Respubliki Bolgarii  
(for Khasymaki). 2. Predsedatel' Soveta po khudozhestvennomu kon-  
struirovaniya Germanskoy Demokraticheskoy Respubliki (for Kel'm).  
3. General'nyy sekretar' Soveta po tekhnicheskoy estetike pri  
Predsedatele Soveta Ministrov Pol'skoy Narodnoy Respubliki (for  
Shidlovskaya). 4. Zamestitel' ministra promyshlennosti tovarov  
shirokogo potrebleniya, zamestitel' predsedatelya Soveta po  
tekhnicheskoy estetike Chekhoslovatskoy Sotsialisticheskoy  
Respubliki (for Gorak). 5. Direktor TSentra khudozhestvennogo  
konstruirovaniya v Zagrebe Sotsialisticheskoy Federativnoy  
Respubliki Yugoslavii (for Rikhter).

KEL'M, M. [Kelr, M.]

Industrial design at the Leipzig Fair of 1965. Tekh. est. 2 no.8:  
28-31 Ag '65. (MIRA 18:9)

1. Predsedatel' Soveta po khudozhestvennomu konstruirovaniya;  
direktor Tsentral'nogo instituta khudozhestvennogo konstruirovaniya  
Germanskoy Demokraticheskoy Respubliki.

MAS'KO, N.Ye., inzh.; KEL'MA, L.Ya., inzh.

A new evaporator cooler for thermal deaerators. Elek. sta. 33  
no.8:71 Ag '62. (MIRA 15:8)  
(Steam power plants)



PETROV, D.G., dotsent, direktor L'vovskogo nauchno issledovatel'skogo instituta perelivaniya krovi.; LYSENKO, E.V.; KEL'MAN, A.A., kandidat meditsinskikh nauk, direktor L'vovskogo oblastnogo onkologicheskogo dispansera.

Plasmotherapy of inflammatory diseases of the female genitalia. Akush.i gin. no.2:45-47 Mr-Ap '53. (MLRA 6:5)

1. L'vovskiy nauchon-issledovatel'skiy institut perelivaniya krovi. 2. L'vovskiy oblastnyy onkologicheskii dispanser. (Gynecology) (Blood--Plasma)

KEL'MAN, A.A., kandidat meditsinskikh nauk [deceased]; KRUKIYER, M.D.  
(L'vov)

Antitoxic function of the liver in cancer of the cervix uteri during radiotherapy. Klin.med. 33 no.4:85 Ap '55. (MLRA 8:7)

1. Iz L'vovskogo oblastnogo onkologicheskogo dispansera (glavnyy vrach - kandidat meditsinskikh nauk A.A.Kel'man)

(ROENTGEN RAYS, effects,

on liver funct., in ther. of cancer of cervix)

(CERVIX, UTERINE, neoplasms,

ther., x-rays, liver funct. in)

(LIVER FUNCTION TESTS, in various diseases,  
cancer of cervix, in x-ray ther.)

KRYLOVSKIY, S.S.; KEL'MAN, A.B.; OSTROVSKIY, A.N.

Firing refractory raw materials in a fluidized bed. Ogneupory 29  
no.1:13-17 '64. (MIRA 17:3)

1. Gosudarstvennyy institut po proyektirovaniyu predpriyatiy po  
proizvodstvu stali.

KEL'MAN, A.B.; PETROVA, Yu.I.; NUZHNYI, N.A.; PERESADA, M.P.

Mixing burner for natural gas. Ogneupory 30 no.8:8-9  
'65. (MIRA 18:8)

1. Nauchno-issledovatel'skiy i proyektnyy institut metallurgicheskoy  
promyshlennosti (for Kel'man, Petrova). 2. Shamotnyy zavod "Krasnaya  
Zvezda" (for Nuzhnyy, Peresada).

VOLOVINSKAYA, V., kand. tekhn. nauk; RUBASHKINA, S.; POLETAYEV, T.;  
KEL'MAN, B.; MERKULOVA, V.

Improving the quality of hams during salting with the use of  
phosphates and sodium ascorbates and glutamates. Mias. ind. SSSR.  
30 no.4:48-50 '59. (MIRA 12:12)

1. Vsesoyuznyy nauchno-issledovatel'skiy institut myasnoy promyshlen-  
nosti.

(Meat, Salt)

VOLOVINSKAYA, V., kand.tekhn.nauk; ~~KEL'MAN~~, B.

Determination of the moisture absorption capacity of meat. Mias.  
ind. SSSR 31 no.6:47-48 '60. (MIRA 13:12)  
(Meat) (Moisture)

*KEL'MAN, B.M.*  
Category: ~~SSSR/Electronics~~ Electronic Optics

H-3

"APPROVED FOR RELEASE: 06/13/2000" CIA-RDP86-00513R000721510013-1

Abs Jour : Ref Zhur - Fizika, No 1, 1957, No 1662

Author : Baranovskiy, S.N., Kaminskiy, D.L., Kel'man, B.M.  
Title : Double Magnetic Slit

Orig Pub : Zh. tekhn. fiziki, 1955, 25, No 11, 1954-1956

Abstract : Description of the construction and of several characteristics of a system consisting of two magnetic slits. Experiments have shown that a double magnetic slit deflects and focuses an electric beam, forming a linear image of a point source.

OSIPOVA, Ye.N.; LIKESH, Yar.; KHEL'MAN, B.M.

Evaluating the performance of fluidisation kilns. Trudy LTI  
no.54:129-134 '59. (MIRA 13:8)  
(Fluidisation--Equipment and supplies)

VOLOVINSKAYA, V.P., kand.tekhn.nauk; KEL'MAN, B.Ya., mladshiy nauchnyy  
sotrudnik

Effect of the technological factors and various surfaces on  
the adhesive properties of the sausage meat. Trudy VNIIMP  
no.14:10-20 '62. (MIRA 16:8)  
(Sausages)



VOLOVINSKAYA, V.P., kand. tekhn. nauk; KEI'MAN, B.Ya., mladshiy nauchnyy  
sotrudnik

Developing the method for determining moisture absorbability  
of meat. Trudy VNIIMP no.11:128-138 '62. (MIRA 18:2)

VOLOVINSKAYA, V.P., kand. tekhn. nauk; KEL'MAN, B.Ya., mladshiy nauchnyy  
sotrudnik

Use of the penetration method for determining the consistency  
of back fat. Trudy VNIIMP no.12:149-156 '62. (MIRA 18:2)

*KEP'MAN, E. I.*

KEP'MAN, E. I.

Ob otvetstvennosti vozdušnogo perevozchika za vred i ubytki.  
(In: Voprosy vozdušnogo prava, v. 1. Moskva, 1927. p. 179-219)  
Title tr.: Liability of air carriers for losses and damages.

DLC: Law Library

SO: Aeronautical Sciences and Aviation in the Soviet Union, Library of  
Congress, 1955.

**KELMAN F. M.**

PROCESSES AND PROPERTIES INDEX

The absorption of nitric oxide by sulfuric acid. K. M. Malin, F. N. Kelman and M. M. Uspenskaya. J. Chem. Ind. (U. S. S. R.) 19, No. 4, 23-7(1938).—NO is absorbed by H<sub>2</sub>SO<sub>4</sub> as an equiv. mixt. with NO<sub>2</sub>, and the presence of excess NO has no effect on the rate of absorption.  
H. M. Leicester

18

MATERIAL INDEX  
CLASSIFICATION

ASB-14A METALLURGICAL LITERATURE CLASSIFICATION

SECONDARY INDEX

DISTRICT

1ST AND 2ND SECTIONS										3RD AND 4TH SECTIONS									
PROCESSES AND PROPERTIES INDEX																			
<p><i>ch</i></p> <p>The desorption of nitrogen oxides from nitrosylsulfuric acid in sulfuric acid. R. M. Malin, F. N. Kel'man, M. M. Mikhovanova and M. M. Uspenskaya. <i>J. Chem. Ind. (U. S. S. R.)</i> 13, No. 6, 24 N(1968). The rate of desorption of <math>\text{NO}_2</math> when a stream of <math>\text{N}_2</math> is passed through the soln. depends on the rate of desorption of nitrosyl-sulfuric acid and the rate of diffusion of <math>\text{NO}_2</math> through the interface. The first factor is the most important under ordinary conditions, but at very high <math>\text{NO}_2</math> concn. the rate of diffusion becomes more important. The presence of a small amt. of <math>\text{HNO}_3</math> greatly increases the rate of desorption.</p> <p>H. M. Leicester</p>																			
<p>ASIA-ELA METALLURGICAL LITERATURE CLASSIFICATION</p>																			
<p>10000 STIMULANT</p>										<p>100000 REF ORG GAT</p>									
<p>100000 REF ORG GAT</p>										<p>100000 REF ORG GAT</p>									

18

CA

Caking of pyrite and its prevention. K. M. Malin, F. N. Kel'man, and A. S. Grigor'eva. *Khim. Prom.* 1947, No. 3, 14-15.—Tests were run to det. the temp. at which pyrite start caking, the loss of S through caking, and means of preventing it. The several kinds of pyrite tested caked at different temps. The caked material contained more FeS than the material that did not cake in all instances but in varying degree. Of the various additives, clay (>15%) and MgO (>5%) reduced caking appreciably. Boiler clinker (>10%) prevented caking entirely. However, the use of additives is undesirable since it interferes with the subsequent use of the pyrite cinders. Mixing of various kinds of pyrite is effective and is recommended if possible. Caking in most instances is prevented if the temp. in any of the hearths exceeds 800°.

M. Hosh

ASM-SLA METALLURGICAL LITERATURE CLASSIFICATION

1947

1948

1949

1950

1951

1952

1953

1954

1955

1956

1957

1958

1959

1960

1961

1962

1963

1964

1965

1966

1967

1968

1969

1970

1971

1972

1973

1974

1975

1976

1977

1978

1979

1980

1981

1982

1983

1984

1985

1986

1987

1988

1989

1990

1991

1992

1993

1994

1995

1996

1997

1998

1999

2000

2001

2002

2003

2004

2005

2006

2007

2008

2009

2010

2011

2012

2013

2014

2015

2016

2017

2018

2019

2020

2021

2022

2023

2024

2025

2026

2027

2028

2029

2030

2031

2032

2033

2034

2035

2036

2037

2038

2039

2040

2041

2042

2043

2044

2045

2046

2047

2048

2049

2050

2051

2052

2053

2054

2055

2056

2057

2058

2059

2060

2061

2062

2063

2064

2065

2066

2067

2068

2069

2070

2071

2072

2073

2074

2075

2076

2077

2078

2079

2080

2081

2082

2083

2084

2085

2086

2087

2088

2089

2090

2091

2092

2093

2094

2095

2096

2097

2098

2099

2100

2101

2102

2103

2104

2105

2106

2107

2108

2109

2110

2111

2112

2113

2114

2115

2116

2117

2118

2119

2120

2121

2122

2123

2124

2125

2126

2127

2128

2129

2130

2131

2132

2133

2134

2135

2136

2137

2138

2139

2140

2141

2142

2143

2144

2145

2146

2147

2148

2149

2150

2151

2152

2153

2154

2155

2156

2157

2158

2159

2160

2161

2162

2163

2164

2165

2166

2167

2168

2169

2170

2171

2172

2173

2174

2175

2176

2177

2178

2179

2180

2181

2182

2183

2184

2185

2186

2187

2188

2189

2190

2191

2192

2193

2194

2195

2196

2197

2198

2199

2200

2201

2202

2203

2204

2205

2206

2207

2208

2209

2210

2211

2212

2213

2214

2215

2216

2217

2218

2219

2220

2221

2222

2223

2224

2225

2226

2227

2228

2229

2230

2231

2232

2233

2234

2235

2236

2237

2238

2239

2240

2241

2242

2243

2244

2245

2246

2247

2248

2249

2250

2251

2252

2253

2254

2255

2256

2257

2258

2259

2260

2261

2262

2263

2264

2265

2266

2267

2268

2269

2270

2271

2272

2273

2274

2275

2276

2277

2278

2279

2280

2281

2282

2283

2284

2285

2286

2287

2288

2289

2290

2291

2292

2293

2294

2295

2296

2297

2298

2299

2300

2301

2302

2303

2304

2305

2306

2307

2308

2309

2310

2311

2312

2313

2314

2315

2316

2317

2318

2319

2320

2321

2322

2323

2324

2325

2326

2327

2328

2329

2330

2331

2332

2333

2334

2335

2336

2337

2338

2339

2340

2341

2342

2343

2344

2345

2346

2347

2348

2349

2350

2351

2352

2353

2354

2355

2356

2357

2358

2359

2360

2361

2362

2363

2364

2365

2366

2367

2368

2369

2370

2371

2372

2373

2374

2375

2376

2377

2378

2379

2380

2381

2382

2383

2384

2385

2386

2387

2388

2389

2390

2391

2392

2393

2394

2395

2396

2397

2398

2399

2400

2401

2402

2403

2404

2405

2406

2407

2408

2409

2410

2411

2412

2413

2414

2415

2416

2417

2418

2419

2420

2421

2422

2423

2424

2425

2426

2427

2428

2429

2430

2431

2432

2433

2434

2435

2436

2437

2438

2439

2440

2441

2442

2443

2444

2445

2446

2447

2448

2449

2450

2451

2452

2453

2454

2455

2456

2457

2458

2459

2460

2461

2462

2463

2464

2465

2466

2467

2468

2469

2470

2471

2472

2473

2474

2475

2476

2477

2478

2479

2480

2481

2482

2483

2484

2485

2486

2487

2488

2489

2490

2491

2492

2493

2494

2495

2496

2497

2498

2499

2500

2501

2502

2503

2504

2505

2506

2507

2508

2509

2510

2511

2512

2513

2514

2515

2516

2517

2518

2519

2520

2521

2522

2523

2524

2525

2526

2527

2528

2529

2530

2531

2532

2533

2534

2535

2536

2537

2538

2539

2540

2541

2542

2543

2544

2545

2546

2547

2548

2549

2550

2551

2552

2553

2554

2555

2556

2557

2558

2559

2560

2561

2562

2563

2564

2565

2566

2567

2568

2569

2570

2571

2572

2573

2574

2575

2576

2577

2578

2579

2580

2581

2582

2583

2584

2585

2586

2587

2588

2589

2590

2591

2592

2593

2594

2595

2596

2597

2598

2599

2600

2601

2602

2603

2604

2605

2606

2607

2608

2609

2610

2611

2612

2613

2614

2615

2616

2617

2618

2619

2620

2621

2622

2623

2624

2625

2626

2627

2628

2629

2630

2631

2632

2633

2634

2635

2636

2637

2638

2639

2640

2641

2642

2643

2644

2645

2646

2647

2648

2649

2650

2651

2652

2653

2654

2655

2656

2657

2658

2659

2660

2661

2662

2663

2664

2665

2666

2667

2668

2669

2670

2671

2672

2673

2674

2675

2676

2677

2678

2679

2680

2681

2682

2683

2684

2685

2686

2687

2688

2689

2690

2691

2692

2693

2694

2695

2696

2697

2698

2699

2700

2701

2702

2703

2704

2705

2706

2707

2708

2709

2710

2711

2712

2713

2714

2715

2716

2717

2718

2719

2720

2721

2722

2723

2724

2725

2726

2727

2728

2729

2730

2731

2732

2733

2734

2735

2736

2737

2738

2739

2740

2741

2742

2743

2744

2745

2746

2747

2748

2749

2750

2751

2752

2753

2754

2755

2756

2757

2758

2759

2760

2761

2762

2763

2764

2765

2766

2767

2768

2769

2770

2771

2772

2773

2774

2775

2776

2777

2778

2779

2780

2781

2782

2783

2784

2785

2786

2787

2788

2789

2790

2791

2792

2793

2794

2795

2796

2797

2798

2799

2800

2801

2802

2803

2804

2805

2806

2807

2808

2809

2810

2811

2812

2813

2814

2815

2816

2817

2818

2819

2820

2821

2822

2823

2824

2825

2826

2827

2828

2829

2830

2831

2832

2833

2834

2835

2836

2837

2838

2839

2840

2841

2842

2843

2844

2845

2846

2847

2848

2849

2850

2851

2852

2853

2854

2855

2856

2857

2858

2859

2860

2861

2862

2863

2864

2865

2866

2867

2868

2869

2870

2871

2872

2873

2874

2875

2876

2877

2878

2879

2880

2881

2882

2883

2884

2885

2886

2887

2888

2889

2890

2891

2892

2893

2894

2895

2896

2897

2898

2899

2900

2901

2902

2903

2904

2905

2906

2907

2908

2909

2910

2911

2912

2913

2914

2915

2916

2917

2918

2919

2920

2921

2922

2923

2924

2925

2926

2927

2928

2929

2930

2931

2932

2933

2934

2935

2936

2937

2938

2939

2940

2941

2942

2943

2944

2945

2946

2947

2948

2949

2950

2951

2952

2953

2954

2955

2956

2957

2958

2959

2960

2961

2962

2963

2964

2965

2966

2967

2968

2969

2970

2971

2972

2973

2974

2975

2976

2977

2978

2979

2980

2981

2982

2983

2984

2985

2986

2987

2988

2989

2990

2991

2992

2993

2994

2995

2996

2997

2998

2999

3000

3001

3002

3003

3004

3005

3006

3007

3008

3009

3010

3011

3012

3013

3014

3015

3016

3017

3018

3019

3020

3021

3022

3023

3024

3025

3026

3027

3028

3029

3030

3031

3032

3033

3034

3035

3036

3037

3038

3039

3040

3041

3042

3043

3044

3045

3046

3047

3048

3049

3050

3051

3052

3053

3054

3055

3056

3057

3058

3059

3060

3061

3062

3063

3064

3065

3066

3067

3068

3069

3070

3071

3072

3073

3074

3075

3076

3077

3078

3079

3080

3081

3082

3083

3084

3085

3086

3087

3088

3089

3090

3091

3092

3093

3094

3095

3096

3097

3098

3099

3100

3101

3102

3103

3104

3105

3106

3107

3108

3109

3110

3111

3112

3113

3114

3115

3116

3117

3118

3119

3120

3121

3122

3123

3124

3125

3126

3127

3128

3129

3130

3131

3132

3133

3134

3135

3136

3137

3138

3139

3140

3141

3142

314

KEL'MAN (and) F. N.

USSR/Pyrites  
Furnaces - Design

"Tendency of Pyrites to Form Clinkers and Methods of Elimination," Malin, Kel'man, Grigor'eva,  
2 pp

"Khimi Promysh" No 3

Discusses susceptibility of various types of ores (Revdinsk, Belevinsk, etc.) , and the process of burning out sulfur in the Karelo-Finnish streaked pyrites at various temperatures. A new type of furnace should be perfected to burn out clinkers at a temperature of about 1100°

PA 10T35

*Sci Res Inst. po udobreniyam i insektofungitsidam*

AMELIN, A.G.; BALEYEV, A.V. [deceased]; BRUTSKUS, Ye.B.; KEL'MAN, F.N.;  
OSHEROVICH, R.Ye.; STEPANOV, M.N.; CHEPYLEVTSKIY, M.L.; CHERNO-  
BAYEVA, M.M.; MIKHAL'CHUK, B.V., redakter; LEONT'YEVA, K.D., re-  
dakter; SHPAK, Ye.G., tekhnicheskij redakter.

[Methods of analyzing and controlling the production of sulfuric  
acid and superphosphates] Metody analiza i kontrolya proizvedstva  
sernei kisloty i superfosfata. Sost. A.G. Amelin i dr. Pod red.  
B.V. Mikhail'chuka. Moskva, Gos. nauchno-tekhn. izd-vo khim. lit-ry,  
1955. 159 p. (MLRA 9:5)

1. Moscow. Nauchnyy institut po udebreniyam i insektofungitsidam.  
(Sulphurec acid) (Phosphates)



AUTHOR: Kel'man, F. N. SOV/32-24-9-8/53

TITLE: An Accelerated Method for the Determination of Selenium in Goods of Sulfuric Acid Production (Uskorennyy metod opredeleniya selena v materialakh sernokislotochnogo proizvodstva)

PERIODICAL: Zavodskaya Laboratoriya, 1958, Vol 24, Nr 9, pp 1061-1064 (USSR)

ABSTRACT: There exist several selenium determination methods, among them that of M. N. Smirnov (Ref 3). The latter has the disadvantage that oxygen is used for the oxidation and that selenium has to be separated from tellurium. The present method is based on the thermal decomposition of the sample in the electric furnace at about 750°, in an air current containing oxides of nitrogen. Selenium is gasified in the dioxide form, is collected and determined iodometrically. The time involved in the analysis is stated to be 1,5 - 2 hours. A diagram of the experimental layout is given, together with a description and the procedure of analysis. The laboratory technicians G. L. Drobinskaya and G. F. Mikheyeva participated in these experiments. Besides  $\text{SeO}_2$ ,  $\text{TeO}_2$  and  $\text{As}_2\text{O}_5$  are volatilized in the thermal decomposition, without, however, interfering with the iodometric

Card 1/2

SOV/32-24-9-8/53

An Accelerated Method for the Determination of Selenium in Goods of Sulfuric Acid Production

selenium determination at a pH = 0,9 - 1,5. In order to prevent a possible volatilization of elementary selenium (without oxidation to  $\text{SeO}_2$ ) it is pointed out that the heating to about 750° should be carried out slowly, so as to facilitate a complete oxidation of selenium. Several tables of analysis results for various materials are given, and evidence is provided for the advantage over other methods of the method described. There are 1 figure, 4 tables, and 6 references, 5 of which are Soviet.

ASSOCIATION: Nauchnyy institut po udobreniyam i insektofungisidam im. Ya. V. Samoylova (Scientific Institute of Fertilizers and Insecto-Fungicides imeni Ya. V. Samoylov)

Card 2/2

KEL'MAN, F.N.; KIRYUSHKINA, M.S.

Improved method for determining arsenic in vanadium catalysts.  
[Trudy] NIUIF no.164:45 '59. (MIRA 15:5)  
(Arsenic--Analysis)

KEL'MAN, Faina Natanovna; BRUTSKUS, Yelena Borisovna; OSHEROVICH,  
Rakhil' Khaymovna. Prinimali uchastiye: GERBURT, Ye.V.;  
MIKHAYL'CHUK, B.V.; SHPAK, Ye.G., tekhn. red.

[Methods of analysis in the control of the production of  
sulfuric acid and phosphorous fertilizers] Metody analiza  
pri kontrole proizvodstva sernoi kisloty i fosfornykh  
udobrenii. Moskva, Goskhimizdat, 1963. 351 p.  
(MIRA 17:2)

KEL'MAN, Faina Natanovna; BRUTSKUS, Yelena Borisovna; OSHEROVICH,  
Rekhal' Khaimovna; MIKHAL'CHUK, B.V., red.; ODERBERG,  
L.N., red.

[Analysis methods in the production control of sulfuric  
acid and phosphorous fertilizers] Metody analiza pri  
kontrole proizvodstva sernoi kisloty i fosfornykh udob-  
renii. Moskva, Khimiia, 1965. 390 p. (MIRA 18:12)

KEELMAN, G.Ya.

Toxicology of paraoxyneozone and "Antox." Kauch. i rez. 24  
no.8:44-45 '65. (MIRA 18:10)

1. Sanitarno-epidemiologicheskaya stantsiya Moskv.

KEL'MAN, G.Ya.

Comparison evaluation of the toxic properties of rubber  
antioxidants. Kauch. i rez. 22 no.12:39-40 D '63. (MIRA 17:9)

1. Sanitarno-epidemiologicheskaya stantsiya, Moskva.

KRIPAK, N.; KEL'MAN, I.; BRAZHNIKOV, V.

Our experience in modernizing production. Mias. ind. SSSR 29  
no.6:8-13 '58. (MIRA 11:12)

1.Semipalatinskiy myasokombinat.  
(Meat industry--Equipment and supplies)



L 45674-66 ENT(m)/T WE  
ACC NR: AP6023622

SOURCE CODE: UR/0318/66/000/004/0012/0015

AUTHOR: Agafonov, A. V.; Osipov, L. N.; Rogov, S. P.; Uzunkoyan, P. N.; Finel'onov, V. P.; Zhandanovskiy, N. B.; Porozhigina, I. Ya.; Kel'man, I. V.; Pisarchik, A. N.; Afanas'yev, V. I.; Khavkin, V. A.; Laz'yan, N. G. 111 116

ORG: All-Union Scientific Research Institute of Petroleum Refining (Vsesoyuznyy nauchno-issledovatel'skiy institut po pererabotke nefti); Novokuybyshev Petroleum Refinery (Novokuybyshevskiy neftepererabatyvayushchiy zavod) 131 116

TITLE: Experience with catalytic hydrocracking of vacuum distillate on the hydrofining assembly of the Novokuybyshev Petroleum Refinery

SOURCE: Neftopererabotka i neftekhimiya, no. 4, 1966, 12-15

TOPIC TAGS: catalytic cracking, petroleum product, gas oil fraction, diesel fuel, gasoline

ABSTRACT: The VNIINP has developed a variant of the process for producing diesel fuel involving one-step hydrocracking of sulfur-containing vacuum distillates on an alumina-cobalt-molybdenum catalyst. The results of laboratory experiments with this variant were successfully applied at the experimental industrial hydrofining assembly of the Novokuybyshev Petroleum Refinery. The operation of the hydrocracking assembly is described. The feed stock for the plant hydrocracking was vacuum gas oil obtained from distillation of sulfur feed stock. Distillation of the hydrogenate produced:

UDC: 665.644.2.048.51665.658.2

Card 1/2

L 45674-66

ACC NR: AP6023622

diesel oil which met all the requirements of GOST 4749-49 for DL grade; a gasoline fraction characterized by a low sulfur content (0.002-0.03), a relatively heavy fractional composition (melting range 120-180°), and a low octane number (42), and is recommended as feed stock for catalytic reforming; the gaseous products methane (49.2 wt. %), ethane (29.4%), propane (17.8%) and butanes (3.65). The residue of the distillation of fuel fractions is recommended as feed stock for catalytic cracking. It is concluded that the hydrocracking of vacuum gas oil on the hydrofining assembly of NKNPZ confirmed the results of work carried out by the VNIINP on pilot plants for the purpose of designing high-capacity units. Orig. art. has: 1 figure and 2 tables.

SUB CODE: 11/ SUBM DATE: none/ ORIG REF: 001/ OTH REF: 003

Card 2/2 fv

KURKO, V., kand. tekhn. nauk; KEL'MAN, L.

Separation of dimethyl esters of pyrogallol and its homologs  
by means of paper chromatography. Mias ind SSSR 34 no. 6:  
50-52 '63. (MIRA 17:5)

1. Vsesoyuznyy nauchno-issledovatel'skiy institut myasnoy  
promyshlennosti.

KURKO, V.I., kand. tekhn. nauk; KEL'MAN, L.F., mladshiy nauchnyy sotrudnik

Chemical nature of smoke components. Trudy VNIIMP no.11:106-118  
(MIRA 18:2)  
(2.

KURKO, V.I., kand. tekhn. nauk; KEL'MAN, L.F., mladshiy nauchnyy sotrudnik

Phenols content of sausage products as indicator of their  
smokiness. Trudy VNIIMP no.12:83-91 '62. (MIRA 18:2)

KEL'MAN L. F.

USSR/Human and Animal Physiology. Metabolism. Nutrition.

Abs Jour: Ref Zhur-Biol., No 12, 1958, 55332.

Author : Belen'kiy, N.G., Krylova, N.N., Chertkov, I.L., Bazarova, K.I.,  
Zuyeva, L.D., Sevost'yanov, B.A., Kel'man, L.F.

Inst : All-Union Academy of Agricultural Sciences.

Title : The Influence of Thermal Treatment on the Assimilation of Meat  
Protein

Orig Pub: Dokl. VASKhNIL, 1957,<sup>22</sup> No 4, 23-29.

Abstract: During a period of 6 days, 26 rats of 180-200 gr body weight each, received daily 10 gr of beef meat with methionine-S<sup>35</sup> proteins. Seven control rats were given raw ground meat. Nine rats were fed ground meat which has been heated in an ultrathermostate at 80° C for one hour, and 10 rats received ground meat heated in an autoclave at 120° C. Two days after the last (6th) feeding, all rats were killed. The proteins were extracted from their plasmas and livers, and their radioactivity was determined. The assimilation of proteins in their natural state as compared to those denaturated by heat

Card : 1/2 A-U Sci Res Inst. 21 Meat Industry

USSR/Human and Animal Physiology. Metabolism. Nutrition.

Abs Jour: Ref Zhur-Biol., No 12, 1958, 55322.

did not show any differences. Thereafter, this investigation was continued on dogs (numbering 8), whereby the nitrogen balance was studied as well. Here, it was established that natural proteins are assimilated somewhat better than denaturalized proteins. Also, it was established that the degree of denaturalization does not exert any specific influence upon protein assimilation.

Card : 2/2

KURKO, V.I., kand.tekhn.nauk; KEL'MAN, L.F., mladshiy nauchnyy sotrudnik

Aromatic properties of phenols, products of thermal decomposition  
of wood. Trudy VNIIMP no.14:36-48 '62. (MIRA 16:8)  
(Meat, Smoked) (Phenols)



KURKO, V.I., kand. tekhn. nauk; KEL'MAN, L.F., mladshiy nauchnyy sotrudnik;  
ROGOV, I.A., kand. tekhn. nauk

Some comparative studies of conventional and electrostatic smoking.  
Trudy VNIIMP no.12:92-103 '62. (MIRA 18:2)

1. Vsesoyuznyy nauchno-issledovatel'skiy institut myasnoy  
promyshlennosti (for Kurko, Kel'man). 2. Moskovskiy  
tekhnologicheskiy institut myasnoy i molochnoy promysh-  
lennosti (for Rogov).

KURKO, V.I., kand. tekhn. nauk; KEL'MAN, L.F., inzh.-khimik;  
MEL'TSER, F.R., inzh.-khimik; KUZNETSOVA, A.A., laborant

Comparative phenol characteristics of smoking preparations  
and uncooked smoked sausage. Trudy VNIIMP no.16:211-220 '64.  
(MIRA 18:11)

ALEKSANDROV, S.N., inzh; KEL'MAN, L.Ya., inzh; PLISAN, I.G., inzh;  
KAMENSKIY, S.K., inzh; RUVIMSKIY, I.M., inzh

Improving the feed-water tubing circuit. Elek.sta 29 no.9:58-64  
S '58. (MIRA 11:11)

1. Pridneprovskaya gosudarstvennaya rayonnaya elektricheskaya  
stantsiya.

(BOILERS)

KEL'MAN, L.Ya., inzh.; MAS'KO, N.Ye.

Operation of ash dumps in electric power plants. Elek. stat.  
35 no.1:11-13 Ja '64. (MIRA 17:6)

KRL'MAN, L.Ya., inzh.

Arrangement for feeding TKZ boilers. Elek.sta. 31 no.7:85 JI  
'60. (MIRA 13:8)

(Boilers)

KEL'MAN, L.Ya., inzh.; MAS'KO, N.Ye., inzh.

Use of shot in the cleaning of a feed water economizer in a system operating on Estonian shale. Elek. sta. 34 no.11:16-20 N '63.  
(MIRA 17:2)

YAKOBSON, A.N., inzh.; TITOV, P.P., inzh.; VERNER, Ye.V., inzh.; KEL'MAN,  
M.M., inzh.

Automatic unit for molding ornamental ceramic tiles. Stroil dor.  
mashinostr. 5 no.3:25-28 Mr '60. (MIRA 13'6)  
(Tiles)

KOZYREV, G.S., dots.; KHEL'MAN, N.I.

Variations ~~in the~~ development of leg muscles in different duck  
breeds. Uch.zap. KHGU 52:245-263 '54. (MIRA 11:11)

1. Kafedra zoologii pozvonochnykh Khar'kovskogo gosudarstvennogo  
universiteta (sav. - prof. I.B. Volchanetskiy).  
(Duck breeds) (Extremities (Anatomy))



L 6956-66 ENT(1)/FCC/EWA(h) CW

ACC NR: AP5026229

SOURCE CODE: UR/0048/65/029/010/1865/1869

AUTHOR: Glikman, L.G.; Kel'man, V.M.; Yakushev, Ye.M.

ORGAN: Institute of Nuclear Physics, Academy of Sciences, KazSSR (Institut yadernoy fiziki Akademii nauk KazSSR)

TITLE: On the electromagnetic mechanism of cosmic ray acceleration /Report, All-Union Conference on Cosmic Ray Physics held at Apatity, 24-31 August 1964/

SOURCE: AN SSSR. Izvestiya. Seriya fizicheskaya. v. 29, no. 10, 1965, 1865-1869

TOPIC TAGS: Primary cosmic ray, particle acceleration, alternating magnetic field, relativistic particle

ABSTRACT: The relativistic equations of motion of a charged particle moving in the plane of antisymmetry of a varying axially symmetric magnetic field are solved for the case when the azimuthal component of the vector potential in the plane of antisymmetry has the form  $f(r/(t - a))/r$ , where  $f$  is an arbitrary function,  $r$  is the distance from the axis,  $t$  is the time, and  $a$  is a constant. Numerical solutions were computed for a field which alternately increases and decreases between finite limits and remains constant for a time at each limit. For the computations it was assumed that the field strength oscillates between  $1.0 \times 10^{-5}$  and  $1.2 \times 10^{-5}$  Oe with a period of  $3.5 \times 10^5$  sec. Some of these solutions are presented graphically and are discussed. The computations show that the ratio of particle energy to field strength is not constant and

Cord 1/2

L 6956-66

ACC NR: AP5026229

that particles can be accelerated to high energies by variable magnetic fields which do not increase indefinitely in strength. Orig. art. has: 19 formulas and 4 figures.

SUB CODE: AA SUBM DATE: 00/--Oct65 ORIG. REF: 006 OTH REF: 000

Card 2/2

L 10664-66 EWT(d)/EWT(1) LIP(c) 00

ACC NR: AP5028313

SOURCE CODZ: UR/0057/65/035/011/1997/2003

AUTHOR: Glikman, L.G.; Kel'man, V.M.; Yakushev, Ye.M.

ORG: none

TITLE: Solution of the nonrelativistic equations of motion for a charged particle in a certain class of varying electromagnetic fields

SOURCE: Zhurnal tekhnicheskoy fiziki, v.35, no. 11, 1965, 1997-2003

TOPIC TAGS: charged particle, motion equation, electromagnetic field, mathematic method

ABSTRACT: The solution of the nonrelativistic equations of motion for a certain class of motions of a charged particle in a certain class of varying electromagnetic fields is reduced to quadratures and eliminations. The electromagnetic fields considered are those that are axially symmetric, have a median plane which is a plane of symmetry for the electric field and a plane of antisymmetry for the magnetic field, and for which the radial and axial components of the vector potential vanish in the median plane (in the gauge in which the scalar potential vanishes) and the azimuthal component of the vector potential in the median plane has the form  $F(r^2/(at^2 + bt + d))/r$ , where  $r$  is the distance from the axis,  $t$  is the time,  $a$ ,  $b$ , and  $d$  are constants, and  $F$  represents an arbitrary function. The motions considered are those in which the particle remains in the median plane. The particular form of the vector

Card 1/2

UDC: 537.533.3

L 10664-66

ACC NR: AP5028313

APPROVED FOR RELEASE: 06/13/2000 CIA-RDP86-00513R000721510013-1  
potential was investigated because it leads simply to an integral of motion. The treatment is different depending on whether the polynomial  $at^2 + bt + d$  does or does not vanish during the motion, and special discussion is required for the case in which the particle passes through the point  $r = 0$ . No applications are suggested for the results obtained. Orig. art. has: 38 formulas.

SUB CODE: 20

SUBM DATE: 12Apr65/

ORIG.REF: 003

OTH REF: 001

Card

2/2 pw

8

Magnetic electron mirror. V. M. Kellman, M. I. Korsunskii and P. P. Lange. *J. Exp. Theoret. Phys.* (U. S. S. R.) 9, 681-4 (1969). There is described a magnetic electron mirror, optically equiv. to a system of cylindrical lenses and a mirror, for directing a nonhomogeneous (polychromatic) electron beam. P. H. Rathmann.

3

Lab. Impact stresses, AS USSR Khaikov



3

*β*-Spectrograph with inhomogeneous magnetic field  
M. Kozminskii, V. Kefman, and B. Petrov. *J. Exptl. Theoret. Phys.* (U.S.S.R.) 16, 301-301 (1911). The *β*-spectrograph capable of exact focusing is discussed. With this *β*-spectrograph a complete focusing of an electron beam in the plane perpendicular to the magnetic field can be obtained within the limits of 40°.  
Galina M. Lebedeff

Physico-Tech Inst, AS USSR

438-114 TOTAL LITERATURE CLASSIFICATION

1ST AND 2ND EDITIONS

PROCESSING AND PROPERTY INDEX

3

CA

$\beta$ -Spectrograph with an inhomogeneous magnetic field.  
M. Korsunskii, V. Kel'man and B. Petrov. (Phys.  
Tech. Inst., Acad. Sci. Ukrainian SSR). *J. Phys.*  
(U.S.S.R.) 9, 7-13 (1945).—See C.A. 39, 2036<sup>h</sup>.

ASACSLA METALLURGICAL LITERATURE CLASSIFICATION

62

KELMAN, V. M.

180T109

USSR/Physics - Spectrography, Beta-Rays Apr 51

"Problem of Beta-Spectrograph Construction Based on  
Analogy With an Optical Spectrograph," V. M. Kelman,  
D. L. Kaminskiy, Leningrad Physicotech Inst, Acad  
Sci USSR

"Zhur Eksper i Teoret Fiz" Vol XXI, No 4, pp 555-561

Authors criticize spectrograph by Klemperer (cf.  
"Phil Mag" 20, 545, 1935) as not similar to opt instr  
and consider its dispersion as zero. Authors design  
new spectrograph analogous to opt and det its dis-  
persion and line width.

LC

180T109



USSR/Physics - Electron Optics

Dec 51

"Motion of Charged Particles in a Magnetic Field of a Linear Current and in the Electric Field of a Cylindrical Condenser," M. Kelman, I. V. Rodnikova, Leningrad Phys Tech Inst, Acad Sci USSR

"Zhur Eksp 1 Teoret Fiz" Vol XXI, No 12, pp 1364-1369

Derives expressions in form of quadratures, detg motion of charged particles in a constant electromagnetic field of a cylindrical condenser with linear conductor along its axis. At certain ratio of field tension and initial velocity of particles such a system

1987103

USSR/Physics - Electron Optics  
(Contd)

Dec 51

has the property to focus bundles of charged particles. Shows that linear charged conductor acts on motion of charged particles with initial velocity perpendicular to conductor as deviating electrooptical element. Submitted 5 Jan 51.

1987103

KELMAN, V. M.

KELMAN, V. M.

USSR/Nuclear Physics - Electron Lens

Apr 52

"Discussion: Modeling the Motion of Charged Particles in Axial-Symmetrical Magnetic Fields,"  
D.L. Kaminskiy, V.M. Kelman

"Zhur Tekh Fiz" Vol XXII, No 4, pp 703-706

Authors criticize article by I.I. Tsukerman (Zhur Tekh Fiz" Vol XXI, 599, 1951) who states that an axial-symmetrical magnetic field scatters electrons similarly to a concave optical lens. Although it may occur in very particular cases, authors consider it impossible to construct a magnetic scattering lens. Received 11 Dec 51.

216T93

USSR/Nuclear Physics - Modeling Motion May 52  
of Particles

"Modeling the Motion of Charged Particles in a  
Two-Dimensional Electric Field Taking Into Ac-  
count the Volumetric Charge," B. V. Bobykin,  
V. M. Kelman, D. L. Kaminskiy

"Zhur Tekh Fiz" Vol XXII, No 5, pp 736-743

In order to find trajectories of charged particles  
in a plane free of space charge, the trajectories  
are considered to be those of balls rolling on a  
rubber membrane. (cf. P. Klymen, Philips Tech

222165

Rev, 2,231, 1937; V. K. Zvorykin et al. "Pro-  
ceedings of IRE" 27,558, 1939). Author modifies  
this method for the case of space charge. Re-  
ceived 14 Feb 52.

222165

KELMAN, V. M.

USSR/Physics - Spectrometers

Card 1/1 Pub. 43 - 10/11

Authors : Kell'man, V. N.; Kaminskiy, D. L.; and Romanov, V. A.

Title : Beta-spectrometer with greater resolving power

Periodical : Izv. AN SSSR. Ser. fiz. 18/1, 148-154, Jan-Feb 1954

Abstract : The construction of a beta-spectrometer of greater resolving power (with symmetrical path of rays) is announced. The spectrometer consists of an electromagnet with screen, copper vacuum-chamber with two copper tubes attached to it, two magnetic lenses, source retainer and recording device. The components of the electrical magnet are described. The current in the coils is directed in such a way that the magnetic current produced by it in the iron yokes are oriented opposite each other. The magnetic current passes through the gap between the upper and lower iron plates of the yoke which also assume the role of poles. Some results obtained with this beta-spectrometer are listed. Two USSR references (1939-1951). Graphs; drawings.

Institution : Academy of Sciences USSR, Physico-Technical Institute

Submitted : December 15, 1953

USSR/Nuclear Physics - Electron magnetic mirrors

Card 1/1 Pub. 43 - 11/11

Authors : Kel'man, V. M., and Lyubimov, K. V.

Title : Similar trajectories of charged particles in magnetic fields

Periodical : Izv. AN SSSR. Ser. fiz. 18/1, 155-160, Jan-Feb 1954

Abstract : Two simple types of electron magnetic mirrors were investigated to determine their suitability in studying the trajectory variations of charged particles in magnetic fields. The magnetic fields of these mirrors were determined by the vector potentials representing homogeneous functions of coordinates with two and three degrees of homogeneity. The form of the trajectory of a charged particle moving in a magnetic field changes during change in the velocity of the particle and depends also upon the intensity of the magnetic field. Three references: 2-USSR and 1-USA (1933-1944). Graphs; drawings.

Institution : Academy of Sciences USSR, Physico-Technical Institute

Submitted : December 15, 1953

USSR/Nuclear Physics - Beta-spectrometers

Card 1/1      Pub. 43 - 4/97

Authors      : Kel'man, V. M.; Kaminskiy, D. L.; and Romanov, V. A.

Title        : A larger prism beta-spectrometer with two magnetic lenses

Periodical   : Izv. AN SSSR, Ser. fiz. 18/2, 209-214, Mar-Apr 1954

Abstract     : The construction and testing of a larger scale prism-type beta-spectrometer with two magnetic lenses for greater resolving and illuminating power is announced. In principle this spectrometer is not different from the spectrometer model described in a previous report; however, its dimensions are larger and it was constructed with greater perfection. The structural and technical characteristics of the prism-type beta-spectrometer are described in detail. Three references: 2 USSR and 1 USA (1939-1954). Graphs; drawings.

Institution   : .....

Submitted    : .....

*NEELMAN, "M."*

USSR/Physics - Charged particle motion

FD-911

Card 1/1 Pub 153-20/26

Author : Kelman, V. M. and Yavor, S. Ya.

Title : Motion of charged particles in a homogeneous magnetic field on which the magnetic field of a linear current and the electric field of a cylindrical condenser are superposed

Periodical : Zhur. tekhn. fiz. 24, 1329-1332, Jul 1954

Abstract : Expressions defining the motion of charged particles in a homogeneous field on which the magnetic field of a linear current and the electric field of a cylindrical condenser are superposed are obtained in the form of quadratures. Numerical integration is carried out in certain particular cases. One reference, by the same author.

Institution : --

Submitted : July 29, 1953

USSR/ Physics - Electron optics

FD-1037

Card 1/1 : Pub. 153 - 8/23

Authors : Kel'man, V. M. Kaminskiy, D. L., and Yavor, S. Ya.

Title : Experimental investigation of cylindrical magnetic electronic lenses

Periodical : Zhur. tekhn. fiz., 24, 1410-1427, Aug - 1954

Abstract : Discuss results of experimental investigation into the electron-optical properties of the magnetic cylindrical lense whose field differs but slightly from the field of two infinite rectilinear oppositely directed currents and also into the systems consisting of two such lenses. Give graphs showing the relation between object position and image for various current strengths. Thanks Diplomat V. P. Vlasenko. Seven references, 2 USSR (N. I. Shtepa, ZhTF, 216, 1952; A. M. Strashkevich, ZhTF, 91, 1940).

Institution : - -

Submitted : 16 March 1954



USSR/Physics - Beta-spectrometer

FD 410

Card 1/1

Author : Kel'man, V. M., Dusayev, G. S., Malkiel', G. S., and Nevodnichiy, N. N.  
Title : Beta-spectrometer with magnetic prism and one magnetic lens  
Periodical : Zhur. eksp. i teor. fiz. 26, 107-108, Jan 1954  
Abstract : Describes the construction and testing of a beta-spectrometer similar to an optical prismatic spectrometer with one lens. Follow the principles of construction discussed by V. M. Kel'man and D. L. Kaminskiy in their work appearing in this journal (Vol 21, 555, 1951)  
Institution : Leningrad Physicotechnical Institute, Acad Sci USSR  
Submitted : November 5, 1951

KEL'MAN, V.M.

YAVOR, S.Ya.; KEL'MAN, V.M., professor

[Magnetic cylindrical lenses] Magnitnye tsilindricheskie linsy.  
Leningrad, 1955. 9 p. (MLRA 9:3)  
(Electrooptics) (Lenses)

*KEL'MAN, V.M.*

Category : USSR/Electronics - Electronic Optics

H-3

Abs Jour : Ref Zhur - Fizika, No 2, 1957, No 4279

Author : Baranovskiy, S.N., Kaminskiy, D.L., Kel'man, V.M.

Title : Investigation of the Electron-Optical Properties of Straight Magnetic Slits.

Orig Pub : Zh. tekhn. fiziki, 1955, 25, No 4, 610-624

Abstract : An investigation was made of the electron-optical properties of many magnetic slit lenses (cylindrical lenses), having various structural dimensions. The constructions of these lenses and their properties are described. A study was made of the qualitative picture of the distribution of the magnetic field in the lens. A qualitative study of the distribution of the field was carried out with the aid of a ballistic galvanometer in three planes, oriented at different angles ( $\varphi = 90^\circ, 180^\circ$ , and  $135^\circ$ ) relative to the surface of the pole pieces and intersecting under the central line of the non-magnetic gap of the lens. It is shown that the distribution obtained in the planes  $\varphi = 90^\circ$  and  $135^\circ$  are in very close agreement with the field of the isolated single conductor, if the current in this conductor is properly chosen.

Card : 1/2

Category : USSR/Electronics - Electronic Optics

H-3

Abs Jour : Ref Zhur - Fizika, No 2, 1957, No 4279

It is thus possible to find the linear current producing a "equivalent" field. Equations for the distribution of the trajectories of the electrons in the field of the isolated linear current were taken from the work by Kel'man and Roknikov (Zh. eksperim. i teoret. fiziki, 1951, 21, 1364). These equations are used to determine the trajectories which start at the peaks of a conical beam and which then fall into the field of the isolated current. It is shown that the field of the isolated current focuses such a beam, forming focal lines.

The authors describe an experimental camera, in which it is possible to study the electron-optical properties of beams of particles, emerging from a gun and entering into the field of the slit lenses of the above construction. The quality of focusing of the beam into a focal line is checked from the image on a fluorescent screen, which is placed on a mount that can be moved in two mutually perpendicular directions, so that the screen can be placed in any previously specified position.

Photographs of the focal line obtained in this manner are given. It is shown that the results of calculation using the "equivalent" linear current and the results of the experiments are close to each other.

Card : 2/2

KEL'MAN, V.M.; LEBEDEV, A.A., akademik, redaktor; SMIRNOVA, A.V.  
tekhnicheskii redaktor.

[Electron optics] Elektronnaia optika. Moskva, Izd-vo Akademii  
nauk SSSR, 1955. 163 p. (MLRA 8:10)  
(Electron optics)

FD-3178

USSR/Physics - Electron Optics

Card 1/1      Pub. 153-8/21

Authors    :   Kel'man, V. M. and Yavor, S. Ya.

Title       :   Investigation of a cylindrical magnetic lens with an iron shell

Periodical:   Zhur. tekhn. fiz., 25, No 8 (August), 1955, 1405-1411

Abstract   :   The authors investigate the electron-optical properties of a jacketed cylindrical magnetic lens encased in iron plates 80 cm long. After a physical description of the apparatus they outline its operational characteristics, expressing the data in graphical form. They give the curve of field distribution, variation in field intensity, dependence of angle of rotation at various lens-to-object distances, as well as other curves expressing various interrelationships among these characteristics.

Submitted   :   March 9, 1955

KEL'MAN, V.M.; KRASHOV, I.F.

Rubber diaphragm technique for determining electron current  
magnitudes in vacuo. Zhur.tekhn.fiz. 25 no.10:1714-1725 S  
'55. (MIRA 9:1)

(Electrons) (Electric discharges through gases)

KRASNOV, I.F.; KEL'MAN, V.M.

Rubber diaphragm technique for solving problems on plane diodes  
with limited emitting surface widths. Zhur.tekh.fiz. 25 no.10:  
1726-1734 S '55. (Diodes) (MLRA 9:1)



Kel'man, V. M.

BARANOVSKIY, S.N.; KAMINSKIY, D.L.; KEL'MAN, V.M.

A double magnetic slit. Zhur.tekh.fiz. 25 no.11:1954-1956 0 '55.  
(Electron optics) (MLRA 9:1)

KEL'MAN, V.M.; ROMANOV, V.A.; METSKHVARISHVILI, R.Ya.

Measurement of the internal conversion coefficients for L- and  
M-subshells of ThC. Dokl. AN SSSR 103 no.4:577-579 Ag'55.  
(MLRA 8:11)

1. Leningradskiy fiziko-tekhnicheskiy institut Akademii nauk  
SSSR. Predstavleno akademikom A.F.Ioffe  
(Thorium--Isotopes) (Spectrometry) (Nuclear shell theory)

**"APPROVED FOR RELEASE: 06/13/2000**

**CIA-RDP86-00513R000721510013-1**

**APPROVED FOR RELEASE: 06/13/2000**

**CIA-RDP86-00513R000721510013-1"**

KEL'MAN, V.M.

Category : USSR/Nuclear Physics - Structure and Properties of Nuclei

C-4

Abs Jour : Ref Zhur - Fizika, No 2, 1957 No 3200

Author : Kel'man, V.M., Metskhvarishvili, R.Ya., Romanov, V.A., Rusinov, L.I.,  
Konoplev, K.A.

Inst : Leningrad Physicotechnical Institute, Academy of Sciences USSR

Title : Determination of the Ratios of the Internal-Conversion Coefficients for  
the Isomeric Transition of  $\text{In}^{114}$ .


Orig Pub : Dokl. AN SSSR, 1956, 107, No 3, 394-397

Abstract : A prism beta spectrometer with a transmission factor .02% and a resolution 0.04% was used to measure the ratios of the conversion coefficients at various L subshells and also at the K, M, and N shells for the 192-kev isomeric transition in  $\text{In}^{114}$ . The ratios are  $L_I: L_{II}: L_{III} = (0.24 \pm 0.01): (1.27 \pm 0.02): 1$ ;  $L/M = 4.4 \pm 0.1$ ;  $M/N = 4.6 \pm 0.2$ ;  $K/L = 1.32 \pm 0.02$ . All data are in good agreement with the type of multipole transition assumed ( $E4$ ).

Card : 1/1

"APPROVED FOR RELEASE: 06/13/2000

CIA-RDP86-00513R000721510013-1



APPROVED FOR RELEASE: 06/13/2000

CIA-RDP86-00513R000721510013-1"

**"APPROVED FOR RELEASE: 06/13/2000**

**CIA-RDP86-00513R000721510013-1**

**APPROVED FOR RELEASE: 06/13/2000**

**CIA-RDP86-00513R000721510013-1"**

**"APPROVED FOR RELEASE: 06/13/2000**

**CIA-RDP86-00513R000721510013-1**

**APPROVED FOR RELEASE: 06/13/2000**

**CIA-RDP86-00513R000721510013-1"**

**"APPROVED FOR RELEASE: 06/13/2000**

**CIA-RDP86-00513R000721510013-1**

**APPROVED FOR RELEASE: 06/13/2000**

**CIA-RDP86-00513R000721510013-1"**



*Kel'man, V.M.*  
AUTHORS

Kel'man, V.M., Utkin, K.G.,  
Loginova, L.N.

57-9-23/40

TITLE

A Simplified Construction of a System Containing  
a Rubber Membrane for the Determination of the  
Trajectories of Charged Particles in the Presence of  
a Space Charge.

(Uproshchennaya konstruktziya ustanovki s rezinovoy  
membranoy dlya opredeleniya trayektoriy zaryazhennykh  
chastits v prisutstvii ob'yemnogo zaryada.)

PERIODICAL

Zhurnal Tekhn. Fiz., 1957, Vol. 27, Nr 9, pp. 2092-2096  
(USSR)

ABSTRACT

In the papers by V.M. Kel'man and I.V. Krasnov,  
Zhurnal Tekhn. Fiz., 1955, Vol. 25, p. 1714, and p.1726  
it was shown that the accuracy with which the trajectories  
of charged particles was determined by means of rubber  
membranes could be considerably increased by giving up  
modelling particle motion by means of a ball rolling  
on a rubber surface, and by using only graphic methods  
for the construction of trajectories. Giving up using  
balls makes it possible to simplify construction con-  
siderably, because in such a case the pressure modelling  
the space charge on the membrane could be brought to

CARD 1/2

AUTHOR: KEL'MAN, V.M., ROMANOV, V.A., MECCHVARIŠVILI, R.JA., PA - 2057  
KOLJUNOV, V.A.  
TITLE: Investigation of Conversion Lines in the  $\beta$  Spectrum of an  $\text{Eu}^{152}$ ,  
 $\text{Eu}^{154}$  Isotopic Mixture. (Issledovanie konversionnykh liniy v  
 $\beta$ -spektre smesi isotopov  $\text{Eu}^{152}$  i  $\text{Eu}^{154}$ , Russian).  
PERIODICAL: Zhurnal Eksperimental'noi i Teoret.Fiziki, 1957, Vol 32, Nr 1,  
pp 39-47 (U.S.S.R.)  
Received: 3 / 1957 Reviewed: 4 / 1957  
ABSTRACT: The authors recorded the lines of the inner conversion on the  
K shells as well as on the L- and M-subshells of the  $\text{Sm}^{152}$  and  
 $\text{Sm}^{154}$  by means of a prism- $\beta$ -spectrometer of great resolving ca-  
pacity and determined the ratios of the conversion coefficients  
at the energies 122 and 123,2 keV of the transitions. The re-  
solving capacity of the prism spectrometer used here was in-  
creased by the following measures: 1) Shielding of the tubes of  
the spectrometer against extraneous magnetic fields by iron  
rings. 2) The straight gap of the registering device was re-  
placed by a slightly curved gap. 3) A certain modification of  
the feeding of the magnet and of the lens of the spectrometer.  
The L-subshells of the  $\text{Sm}^{152}$  and  $\text{Gd}^{154}$ : A diagram demonstrates  
the sphere of the  $\beta$ -spectrum of a  $\beta$ -spectrum of  $\text{Eu}^{152}$  and  $\text{Eu}^{154}$

Card 1/3

Investigation of Conversion Lines in the  $\beta$  Spectrum of an  
 $\text{Eu}^{152}$ ,  $\text{Eu}^{154}$  Isotopic Mixture.

with those lines that were produced by the electrons of the  
inner conversion of the  $\gamma$ -rays with the energies 122 and  
123,2 keV on the subshells of the  $\text{Sm}^{152}$  and  $\text{Gd}^{154}$ . The condi-  
tions under which the lines were obtained as well as the be-  
havior of the lines are discussed. The theoretical ratios of the  
conversion coefficients agree with the here measured ratios.

The M-subshells of samarium: A further diagram demonstrates  
the sphere of the  $\beta$ -spectrum with the lines which are pro-  
duced by the inner conversion on the M-subshell of samarium.  
The authors found the following ratio of the coefficients of  
the conversion on the M-subshells:  $M_I:M_{II}:M_{III} = 1:(3,4\pm 0,1):$   
 $:(3,3\pm 0,2)$ . This corresponds to the conclusions from the  
approximated calculations of the relative conversion coeffi-  
cients. Furthermore  $L/M = 4,5\pm 0,1$  ( $L=L_I+L_{II}+L_{III}$ ;  
 $M = M_I+M_{II}+M_{III}$ ) was found.

Card 2/3

AUTHORS: Kel'man, V.M., Matskhvarishvili, M.Ia., Romanov, V.A. 56-3-6/59  
Tuchkevich, V.V.,

TITLE: The Investigation of Conversion Lines in the  $\beta$ -Spectrum of  $\text{Ir}^{192}$ .  
(Issledovaniye konversionnykh liniy v  $\beta$ -spektre  $\text{Ir}^{192}$ )

PERIODICAL: Zhurnal Eksperim. i Teoret. Fiziki, 1957, Vol. 33, Nr 3, pp.588-594  
(USSR)

ABSTRACT: With the help of a prism- $\beta$ -spectrometer (resolving of 0,04 %) the  
conversion coefficients and the multipole order of the following  
 $\gamma$ - lines were determined:

E $\gamma$ in KeV	K/L	K/M	multipole order
136,3			(80 $\pm$ 1)% E2 + (20 $\pm$ 1)% M1
201,3	1,85 $\pm$ 0,04		(86 $\pm$ 2)% E2 + (14 $\pm$ 2)% M1
205,8	1,83 $\pm$ 0,04		E2
295 ,8	2,35 $\pm$ 0,04	8,9 $\pm$ 0.2	E2
308,5	2,38 $\pm$ 0,02	9,5 $\pm$ 0,2	(97 $\pm$ 2)% E2 + (3 $\pm$ 2)% M1
316,5	2,22 $\pm$ 0,02	9,3 $\pm$ 0,2	E2
468,0	3,0 $\pm$ 0,1	10,2 $\pm$ 0,2	E2
604,5	4,7 $\pm$ 0,1		(88 $\pm$ 2)% E2 + (12 $\pm$ 2)% M1

Card 1/2

The Investigation of Conversion Lines in the  $\beta$ - Spectrum of  $\text{Ir}^{192}$  56-3-6/59

APPROVED FOR RELEASE: 06/13/2000 CIA-RDP86-00513R000721510013-1"

ASSOCIATION: Leningrad Physicoal-Technical Institute AN USSR  
(Leningradskiy fiziko-tekhnicheskiy institut Akademii nauk SSSR)

SUBMITTED: March 18, 1957

AVAILABLE: Library of Congress

Card 2/2

*KEL' MAN, V.M.*

AUTHOR: DOLMATOVA, K.A., KEL'MAN, V.M. 20-6-16/59  
TITLE: A Longitudinal  $\beta$ -Spectrometer with Compensated Spherical Aberration.  
(Prodol'nyy  $\beta$ -spektrometr s kompensirovannoy sfericheskoy aberratsiyey. Russian).  
PERIODICAL: Doklady Akademii Nauk SSSR, 1957, Vol 113, Nr 6, pp 1244 - 1247 (U.S.S.R.)  
ABSTRACT: The spherical aberration described in the paper under review is compensated by a transverse magnetic field of the field strength  $H = H_0/r$ . The additional F is generated by a winding which concentrates in an annular focus the ions which fly out of the source in a wide solid angle. No difficulty is encountered in computing the electrons paths in the range of the homogeneous field. Also in the range where the homogeneous magnetic field is superposed with a field of the field strength  $H = H_0/r$ , the differential equations of the motion of the electrons are reduced to quadratures. The present paper contains the revalent formulae for the differential equations and for their solutions. The compensating field was applied in the neighborhood of the apex of the orbit. With the aid of these formulae a form was found for the boundary of the inhomogeneous field which guarantees an annular focus of aberration (if a point source is used). This focus coincides with the linear annular focus which is formed

Card 1/3

A Longitudinal  $\beta$ -Spectrometer with Compensated Spherical Aberration.

by an infinitely small bundle of the ions flying away under an angle of  $30^\circ$ .

The longitudinal field was generated by a coil of a length of 110 cm and of an internal diameter of 33 cm; this coil was wound on a copper tube. At the same time, this copper tube also served as chamber of the spectrometer. Additional improvements in the homogeneity of the field were achieved by the use of correcting coils. Almost everywhere the form of the coils was in agreement with the computed form of the boundary of the field. The radioactive source was glued to a fastener which was introduced into the vacuum chamber through a Wilson compression /?/ and through a vacuum tap. A counter (G-M5) with a window of a diameter of 90 mm served as detector. Employment of a compensating field reduces the width of the annular projection by about 2.5 times as compared to the width of the projection in a homogeneous field. Further investigations were conducted with the aid of a radioactive source (active  $Ti^{48}$ -precipitation) with the dimensions of 1 X 1 mm. A brief discussion of the results is given. (4 reproductions).

Card 2/3

AUTHOR: Kel'man, V. M., Doctor of Physical- and Mathematical Sciences SOV/30-58-7-13/49

TITLE: New Magnetic Beta-Spectrometer (Novyy magnitnyy beta-spektrometr)

PERIODICAL: Vestnik Akademii nauk SSSR, 1958, Nr 7, pp. 75 - 78 (USSR)

ABSTRACT: The magnetic prism deflecting the electron beam, must be considered as the most important part of this spectrometer which was developed in the Physical-Technical Institute. The parallel electron beam striking the deflection magnet, must remain parallel also after its deflection. This condition, however, can only be satisfied if a deflecting magnet as represented in figure 1, is used. The construction of the  $\beta$ -spectrometer is given in figure 2 and is then described in detail. The conversion line of  $\gamma$ -quanta of different energies is given in figure 3, being a section of the spectrum of conversion electrons which are produced in the decay of  $\text{Ir}^{192}$ . The conversion lines of the I and Ia active sediment  $\text{RdTh}$  are given in figure 4. A further perfection of the new  $\beta$ -spectrometer is to be achieved by an improvement of the deflection magnet, the electronic lenses and by shielding

Card 1/2

New Magnetic Beta-Spectrometer

SOV/30-58-7-13/49

the electrons against the action of random magnetic fields. It is further suggested to use magnetic material of higher quality and to pay more attention to the stabilization of the magnetic field. This kind of work is carried out at present in the Physico-Technical Institute. There are 4 figures.

Card 2/2

AUTHORS: ~~Kellman, V. M.~~, Peregud, B. P.,  
Dolmatova, K. A.

57-28-5-26/36

TITLE: Accelerators With a Radially Growing Leading Field and Additional Electron Optical Elements for Securing the Vertical Focussing of the Beam (Uskoriteli s radial'no narastayushchim vedushchim polem idopolnitel'nyimi elektronnoopticheskimi elementami, obespechivayushchimi vertikal'nuyu fokusirovku puchka)

PERIODICAL: Zhurnal Tekhnicheskoy Fiziki, 1958, Vol. 28, Nr 5, pp. 1056-1064 (USSR)

ABSTRACT: The application of a radially decreasing field in modern weakly focussing accelerators is determined by the necessity of a vertical focussing of the beam of the accelerated particles. The new possibilities, which have been proposed from various sides (references 1-8) immediately attracted the interest of researchers. Recently, numerous experimental and theoretical investigations were conducted dealing with the application of these proposals in different types of accelerators (references 9-22). All these methods have the following in common: 1) The vertical field component is not constant in the middle plane and periodi-

Card 1/4

Accelerators With a Radially Growing Leading Field and . 57-28-5-26/36  
Additional Electron Optical Elements for Securing the Vertical Focussing  
of the Beam

cally changes its value, or, with respect to the azimuth, even its direction. 2) The functions of the leading and of the focussing field are performed by one and the same field, which only formally can be regarded as a superposition of two fields. This field, however, is created only by one magnetic system; 3) The magnet poles must possess an accurately worked, complicated profile (method by Thomas and the spiral-sector variant) or the field must be created by a great number of accurately placed sector magnets. A series of shortcomings attached to the new accelerator constructions are a result of these circumstances. The authors propose another method. The focalization is effected by supplementary electron optical elements: with cylindrical magnetlenses or magnet gaps. The method guarantees the stability of the radial as well as of the vertical betatron oscillations and can be employed for the construction of circular accelerators of different types. In this paper the possible constructional variants of the focussing system are drawn into consideration and the electron model is described. The peculiarities of the proposed method differentiating it from earlier

Card 2/4



Accelerators With a Radially Growing Leading Field and Additional Electron Optical Elements for Securing the Vertical Focussing of the Beam 57-28-5-26/36

ones, are as follows: 1) A separation of the functions of focalization and of leading the beam between two elements - the magnet and the focussing system. This guarantees the free choice of the shape of the leading field and facilitates its creation. As a result of the separation a facilitated leading of the beam and a slackening of the restrictions imposed upon the production and the mounting of the constructional nodes of the accelerator can appear. This is the case in particular, if small adjustments and a flexibility of the elements of the focussing system during the mounting of the accelerator are provided for. 2) The comparatively low weight of the electromagnet creating the leading field in comparison to the weight necessary in earlier methods. This is connected with the fact that the magnetic circuit of the focussing system is not closed by the yoke of this magnet. 3) An increase of the copper weight and of the necessary power. 4) A more simple construction of the electromagnet consisting of the possibility of employing a closed ring magnet with a low number of magnetizing coils and no sector magnet. An electronic simulator was built for experimental exa-

Card 3/4

Accelerators With a Radially Growing Leading Field and 57-28-5-26/36  
Additional Electron Optical Elements for Securing the Vertical Focussing  
of the Beam

mination. A schematic cross section of this model is shown in figure 2. At present the model is prepared for experiment. The authors thank G.A. Grinberg, Yu.V. Vandakurov, D.G. Alkhazov and D.M. Kaminker. There are 3 figures and 29 references, 10 of which are Soviet.

ASSOCIATION: Fiziko-tekhnicheskii institut AN SSSR, Leningrad (Leningrad, Physical-Technical Institute, AS USSR)

SUBMITTED: July 11, 1957

1. Particle accelerators--Design 2. Particle beams--Focusing

Card 4/4

KUL'MAN, V.M., doktor fiziko-mat. nauk

New magnetic beta-spectrometer. Vest. AN SSSR 28 no. 7:75-78  
J1 '58. (MIRA 11:7)

(Spectrometer)

21(9)

SOV/56-35-5-7/56

AUTHORS:

Kel'man, V. M., Kolyunov, V. A., Karpov, M. V.

TITLE:

The Application of Magnetic Slits for the Creation of Circular Trajectories of Charged Particles (Primeneniye magnitnykh shcheley dlya formirovaniya krugovykh trayektoriy zaryazhennykh chastits)

PERIODICAL:

Zhurnal eksperimental'noy i teoreticheskoy fiziki, 1958, Vol 35, Nr 5, pp 1113-1115 (USSR)

ABSTRACT:

The authors of the present paper investigated an electron-optical system consisting of magnetic slits with a magnetic field increasing rapidly in the direction of the periphery but not leading to defocusing in a vertical direction, which bends the trajectories of charged particles, rendering them nearly circular. The vector potential in point P of this system has the form

$$A_z = -\frac{I}{c} \ln \frac{(r/a)^{2n} - 2(r/a)^n \cos n\varphi + 1}{(r/a)^{2n} + 2(r/a)^n \cos n\varphi + 1}; \quad A_r = A_\varphi = 0, \quad A = A_z.$$

(I = current in every conductor, r = distance between the axis of the system O and P, a = distance from O to conductor,

Card 1/3

SOV/56-35-5-7/56

The Application of Magnetic Slits for the Creation of Circular Trajectories of Charged Particles

$2n$  = the number of conductors, and  $\varphi$  = the polar angle). For the momentum it holds that  $P = \partial L / \partial \dot{z} = m\dot{z} + eA/c = m\dot{z}_0 + eA_0/c = \text{const}$ ; the Lagrangian  $L = \frac{m}{2} (\dot{r}^2 + \dot{\varphi}^2 + \dot{z}^2) + eA\dot{z}/c$ . Scheme

and schematic drawing of such a system which can be used in an accelerator with a constant guiding field are given (Figs 1, 2). The experiments carried out with this device are described. The device consisted of 16 poles arranged in a circle and having 200 windings each; the distance between the gun and the edge of the poles  $\sim 4.5$  cm. at radial oscillation of the order of 2 cm and vertical oscillations  $\sim 5$  cm. The electron energy amounted to 3 keV (5 - 10 A). The amperage depended in a high degree on the distance between gun and pole. The phenomenon had the shape of a slightly curved band of 1 - 3 mm breadth and 10 - 20 mm height. An arrangement consisting of 32 poles gave similar results. There are 2 figures and 4 Soviet references.

Card 2/3

SOV/56-35-5-7/56

The Application of Magnetic Slits for the Creation of Circular Trajectories  
of Charged Particles

ASSOCIATION: Leningradskiy fiziko-tekhnicheskii institut Akademii nauk SSSR  
(Leningrad Physico-Technical Institute of the Academy of  
Sciences, USSR)

SUBMITTED: May 16, 1958

Card 3/3

21(8)

SOV/56-35-5-51/56

AUTHORS:

Kel'man, V. M., Metekhvashvili, R. Ya., Proobrazhanskiy, B.K.,  
~~Romanov, V. A.,~~ Tushkevich, V. V.

TITLE:

The Investigation of the Spectrum of Conversion Electrons of  
the Isotopes of Lutetium With Neutron Deficit (Issledovaniye  
spektra konversionnykh elektronov neytronodefitsitnykh  
izotopov lyutetsiya)

PERIODICAL:

Zhurnal eksperimental'noy i teoreticheskoy fiziki, 1958,  
Vol 35, Nr 5, pp 1309-1310 (USSR)

ABSTRACT:

The investigation of the radiation of greatly deformed nuclei  
furnishes material for the further development of the collective  
nuclear model. It is just from this point of view that the  
isotopes of lutetium are of interest. Recently several papers  
(Refs 1-4) have been published which deal with lutetium  
isotopes with neutron deficit, but the data given by these  
papers do not convey a clear idea of the decay of these iso-  
topes. Additional investigations are therefore necessary. The  
authors of the present paper investigated the conversion spec-  
trum of the isotopes of a lutetium fraction, which had been  
separated from a tantalum target irradiated with fast (660 MeV)  
protons. The method employed for separation has already been

Card 1/3

SOV/56-35-5-51/56

The Investigation of the Spectrum of Conversion Electrons of the Isotopes of Lutetium With Neutron Deficit

described (Ref 5). Measurements were carried out by means of a prism- $\beta$ -spectrometer and by means of a double-focusing spectrometer. The spectrum of the conversion electrons consists of many lines, which belong to  $\text{Lu}^{169}$  (half-life  $\sim 1.5$  days),  $\text{Lu}^{170}$  ( $\sim 2$  days),  $\text{Lu}^{171}$  ( $\sim 8$  days),  $\text{Lu}^{172}$  ( $\sim 6.7$  days),  $\text{Lu}^{173}$  ( $\sim 200$  days). Belonging of lines to the various corresponding isotopes was determined from the half-life. A table gives the energies of  $\gamma$ -transitions the conversion lines of which decrease with the period  $\sim 1.5$  to 2 days. The second table contains the energies of the  $\gamma$ -transitions with the period 6.7 to 8 days. The energy of these transitions was determined from the energy of K- and L-conversion lines. There are 2 tables and 6 references, 4 of which are Soviet.

ASSOCIATION: Leningradskiy fiziko-tekhnicheskoy institut Akademii nauk SSSR (Leningrad Physico-Technical Institute of the Academy of Sciences, USSR)

Card 2/3



PHASE I BOOK EXPLOITATION

SOV/3514

Kel'man, V.M., and S.Ya. Yavor

Elektronnaya optika (Electron Optics) Moscow, Izd-vo AN SSSR, 1959. 372 p.  
3,000 copies printed.

Sponsoring Agency: Akademiya nauk SSSR. Fiziko-tekhnicheskii institut.

Ed.: L.A. Artsimovich, Academician; Ed. of Publishing House: Yu.K. Imshenetskiy;  
Tech. Ed.: A.V. Smirnova.

PURPOSE: The book is intended for students of electron optics.

COVERAGE: The book deals primarily with geometrical electron optics and does not discuss wave properties of electrons. In addition to the theory of focusing particle beams in fields with symmetry of rotation, the theory of focusing in electromagnetic fields with arbitrary space distribution is presented. Cylindrical electron lenses and various kinds of deflection systems are discussed in detail. Much space is devoted to calculations of the motion of charged particles. The last two chapters are devoted to applications of electron optics. No personalities are mentioned. There are 277 references, 119 Soviet (17 are translations), 88 English, 48 German, 13 French, 3 Chinese, and 6 Scandinavian.

Card 1/6

Electron Optics

SOV/3514

TABLE OF CONTENTS:

Introduction	3
Ch. I. General Methods of Solving Problems of the Motion of Charged Particles in Electric and Magnetic Fields	5
1. Motion equations in the Newtonian form	5
2. Lagrange equations	18
3. Hamilton-Jacobi equation	28
4. Analogy between the motion of charged particles in electromagnetic fields and propagation of light in transparent media	35
Ch. II. Approximate Methods of Determining Trajectories of Charged Particles in Electric and Magnetic Fields	42
1. Numerical methods of integrating differential motion equations	42
2. Graphical methods of constructing trajectories	52
3. Modeling of fields and trajectories of charged particles	60
Ch. III. Focusing a Beam of Charged Particles	82
1. G.A. Grinberg's general theory of the focusing action of electric and magnetic fields	82
Card 2/6	

Electron Optics

SOV/3514

2. Focusing in an electric field	94
3. Individual cases of the focusing of charged particles	98
Ch. IV. Electron-Optical Properties of Fields With Rotational Symmetry	107
1. Fields with rotational symmetry	107
2. Motion of electrons in fields with rotational symmetry	113
3. Basic equations of electron optics	115
4. Obtaining images in fields with rotational symmetry	120
Ch. V. General Properties of Electron Lenses	126
1. Classification of electron lenses	126
2. Formation of images in electron lenses	128
3. Approximate methods of calculating trajectories in electron lenses	137
4. Experimental methods of determining the location of cardinal points of electron lenses	141
5. Thin lenses	145
6. Object in the lens field	148
Ch. VI. Electrostatic Lenses	150
1. Diaphragm with round opening	150
2. Immersion lens	155
Card 3/6	

Electron Optics

SOV/3514

3. Unipotential lens	161
4. Cathode lens (immersion objective)	165
5. Electrostatic electron mirrors	172
Ch. VII. Magnetic Lenses	178
1. Long magnetic lens	178
2. Short magnetic lens	180
3. Accurate integration of trajectory equations for certain types of magnetic lenses	185
4. Experimental determination of electron lens parameters	191
Ch. VIII. Aberrations of Electron Lenses	196
1. Calculation of geometric aberrations by the method of the eikonal function (characteristic Hamilton function)	196
2. Characteristic of individual types of geometric aberration	207
3. Calculation of coefficients of geometric aberration	216
4. Spherical aberration	219
5. Chromatic aberration	228
6. Other kinds of aberration	237

Card 4/6